

Case Report

Laparoscopic repair of Amyand hernia – A case report with review of literature

Prachi Praveen Agrawal¹, Abhijit S Joshi²From, ¹DNB 2nd year Resident, ²Consultant, Department of General and Advanced Laparoscopic Surgery, Dr. L. H. Hiranandani Hospital, Powai, Mumbai

ABSTRACT

Amyand's hernia is a rare condition in which the appendix becomes incarcerated within an inguinal hernia. Although this type of hernia is uncommon, it can lead to serious complications such as strangulation and perforation when the appendix becomes trapped. Incarceration of the appendix is most frequently observed in inguinal and femoral hernias, but it can also occur, though less commonly, in incisional and umbilical hernias. This case report describes an 83-year-old male, who was diagnosed with a right inguinal hernia and underwent laparoscopic hernia repair. Intraoperatively, a right-sided Amyand's hernia was identified. Since the appendix was not inflamed, it was reduced and a mesh plasty was performed to repair the hernia. The patient had an uneventful recovery.

Key words: Appendicitis, Hernia, Incarceration, Inguinal, Laparoscopy, Mesh plasty

Amyand's hernia is a rare surgical finding where the vermiform appendix is incarcerated within an inguinal hernia sac. It often presents with nonspecific symptoms and is typically diagnosed intraoperatively [1]. This case report discusses the laparoscopic management of a right-sided Amyand's hernia in an elderly male, emphasizing the role of intraoperative vigilance and appropriate surgical decision-making.

CASE REPORT

An 83-year-old male presented to the surgical outpatients department (OPD) with a complaint of swelling in the right groin region, which became more prominent on straining, with occasional discomfort since 20 days. He had no known comorbidities. He had a history of undergoing open surgical repair of an umbilical hernia 2 years back. On examination, the right inguinal swelling was reducible and exhibited a positive expansile cough impulse; suggestive of right inguinal hernia. The horizontal infra-umbilical scar of the previous umbilical hernia surgery was noted. The patient was then planned for elective laparoscopic right inguinal hernia repair under general anaesthesia. Though the totally extraperitoneal (TEP) is the authors' preferred approach, in view of the infraumbilical scar and previous mesh placement and anticipated adhesions due to the same, it was decided to adopt

the trans-abdominal pre-peritoneal (TAPP) approach. Intraoperatively, the right-sided indirect inguinal hernia was found to contain the appendix and mesoappendix within the hernia sac, thereby confirming an Amyand's hernia (Type 1).




Fig 1. A) Indirect RIH (blue arrow) with appendix as its content (yellow arrow), red 'C' shows intraabdominal caecum, B) Appendix being reduced, C) Reflection of peritoneal flap during TAPP (blue arrow), D) Development of extraperitoneal space (yellow asterisk) by blunt dissection (blue arrow)

Access this article online

Received – 10th May 2025
Initial Review – 14th May 2025
Accepted – 10th June 2025

DOI: 10.32677/ejms.v10i3.7616

Quick Response code



Correspondence to: Dr. Prachi Praveen Agrawal, Department of General and Advanced Laparoscopic Surgery, Dr. L. H. Hiranandani Hospital, Powai, Mumbai.

Email: prachiag1996@gmail.com

©2025 Creative Commons Attribution-Non Commercial 4.0 International License (CC BY-NC-ND 4.0).




Fig 2. A) Further development of extraperitoneal space (yellow asterisks) by blunt dissection (blue arrow) anterior to the reflected peritoneal flap (red arrow), B) Division of fibrotic attachment of mesoappendix to indirect ring (blue arrow), C) Fibrotic attachment (blue asterisk) completely divided & appendix reduced to its normal position in right iliac fossa (yellow arrow), D) Right indirect sac (yellow asterisk) being skeletonised (blue arrow)




Fig 3. A) Division of skeletonised right indirect sac (blue arrow), B) Sac completely divided & pushed proximally (blue asterisk), C) Prolene mesh (blue asterisks) placed optimally & tacked to Cooper's ligament (yellow arrow), D) Previously reflected peritoneal flap re-positioned and tacked to parietes so as to extra-peritonealise the mesh (yellow arrow)

No signs of perforation or abscess were noted (Fig 1A). As the appendix was not inflamed, it was simply reduced (Fig 1B). The peritoneal flap was reflected and extra-peritoneal space developed as part of the hernia repair (Fig 1C, D & 2A). The fibrotic attachment of the meso-appendix to the hernia ring was divided and the appendix was repositioned back to the right iliac fossa (Fig 2 B, C). The indirect sac was then skeletonised and divided (Fig 2D & 3A, B). A standard

meshplasty was performed, using a 12 X 15 cm Prolene mesh and a Protack hernia tacker (Fig 3C). The peritoneal flap was then repositioned back so as to extra-peritonealise the mesh (Fig 3D). The post-operative course was uneventful. During his post-operative day 10 OPD follow-up visit, all his operative wounds had healed well. At the time of writing this paper, a telephonic interview was conducted with the patient. Twelve months after his surgery, he continues to be asymptomatic.

DISCUSSION

The condition is named after Claudius Amyand, who, on December 6, 1735, performed the first successful appendectomy on an 11-year-old boy presenting with a right inguinal hernia. During the surgery, Amyand discovered a pin lodged in the appendix, which was surrounded by stones. The appendix was located within the inguinal hernia sac. Consequently, Amyand's hernia (AH) refers to an incarcerated hernia that contains the appendix, which can either be inflamed or normal [2]. It has been documented in individuals aged from 3 weeks to 92 years, frequently diagnosed in children, being three times more likely than in adults, primarily due to the patency of the processus vaginalis in children [2]. A recent systematic review indicated a higher prevalence of AH in men (87.6%) [3].

The incidence of AH is estimated to be around 0.4%-1% of all inguinal hernias [4]. An incarcerated appendix in an inguinal hernia may become inflamed, infected, or perforated. However, in some cases, the appendix remains completely healthy despite being incarcerated, as seen in our case. Appendicitis within an AH hernia constitutes 0.1% of all cases [4]. The occurrence of a perforated appendix within an inguinal hernia is also rare, accounting for 0.1% of appendicitis cases [4]. In females, AH-related appendicitis is often observed in postmenopausal women [4]. The mortality rate of AH hernia ranges from 14% to 30%, primarily due to sepsis spreading through the peritoneum [4]. Amyand's hernia can involve either direct or indirect inguinal hernias. Diagnosing AH before surgery is challenging, and it is often discovered incidentally during surgery.

Patients commonly present with sudden-onset pain in the epigastric or periumbilical area, localized tenderness in the right lower quadrant, and a tender irreducible mass in the inguinal or inguino-scrotal region. The presence of acute appendicitis in AH remains a controversial subject. It is typically caused by extra-luminal obstruction due to pressure at the hernia neck rather than an intraluminal obstruction [5]. The appendix becomes more vulnerable to injury and is eventually trapped by adhesions when it enters the hernia sac. Increased intra-abdominal pressure, such as muscle contractions, can compress the appendix, causing inflammation and potentially disrupting its blood supply, leading to further inflammation and bacterial growth.

Amyand's hernia is most often found on the right side due to the anatomical position of the appendix.

However, left-sided cases have been reported, potentially linked to conditions such as situs inversus, intestinal malrotation, or a mobile cecum [5]. In these cases, an appendectomy is performed if the cecum's base is accessible through the hernial sac. If it is not, an open appendectomy may be performed through a low-midline laparotomy incision. Some suggest performing an appendectomy even in cases of left-sided AH, even if the appendix appears normal, as any future appendicitis could present atypically and delay diagnosis.

Preoperative diagnosis of AH is rare, with most diagnoses made during surgery. Abdominal exams, physical findings, lab results, and imaging studies are not always helpful in differentiating AH from other conditions. When a normal appendix is discovered during surgery, it often does not present with symptoms. However, an inflamed appendix trapped within the inguinal canal can lead to incarcerated hernias, faecal spillage, and sepsis [3]. Common imaging techniques, such as ultrasonography and CT scans, can assist in diagnosing AH. CT is frequently used to evaluate acute abdominal conditions and abdominal hernias.

However, inguinal hernias are typically diagnosed clinically, and imaging is often ordered to exclude other, more serious conditions. The accuracy of ultrasound as a diagnostic tool is heavily dependent on the operator's skill, making it relatively unreliable [4]. The treatment of AH depends on the intraoperative findings. Differential diagnoses can include orchitis, testicular torsion, ovarian inflammation, and bowel

inflammation. Laparoscopic appendectomy has become a preferred method of surgery, particularly due to its benefits in reducing the risk of surgical site infections. If the appendix is inflamed, appendectomy is recommended without the use of mesh for hernia repair [5].

Losanoff and Basson's classification system helps guide the surgical treatment of different types of AH. Type 1 consists of a normal appendix within the inguinal hernia. Type 2 refers to acute appendicitis within the hernia sac, but without abdominal sepsis. Type 3 is Acute appendicitis with peritoneal or abdominal wall sepsis. Type 4 is acute appendicitis with other intra-abdominal pathology. This type is related to other findings, which may be incidental to the hernia. According to this system, mesh placement is not recommended in cases of appendicitis (Types 2–4). Patients with any form of appendicitis should undergo an appendectomy, followed by natural tissue repair of the hernia. If appendicitis is not present, the appendix should be reduced back into the abdomen, and mesh can be used to repair the inguinal hernia (Type 1) [6].

To summarise, AH is a rare form of inguinal hernia in which the appendix is incarcerated within the hernia sac. Its diagnosis remains challenging due to its low incidence, vague clinical presentation, and unclear imaging findings. As such, surgery often serves both diagnostic and therapeutic purposes. The appendix may remain healthy or become inflamed in these cases. A review of the literature on reported cases of AH over the last 5 years is summarised (Table 1). A list of named external herniae with interesting rare contents &/or anatomy including AH, is summarised (Table 2).

Table 1: A review of recent literature on Amyand Hernia over the last 5 years

Authors	Pre-operative diagnosis	Time of diagnosis / Condition of appendix	Mode of surgery (open/lap; mesh - Y or N; appendectomy – Y or N)
Hina Khalid et al [6]	Right indirect inguinal hernia	Intraoperative / non inflamed (Type 1)	Open / Y / Y
Kasra Hatampour et al [3]	Right indirect inguinal hernia	Intraoperative / Inflamed (Type 2)	Open / Y / N
Shu-Yu Wu et al [7]	Right direct inguinal hernia with an incarcerated appendix	Pre-operative (CT scan) / Inflamed (Type 2)	Lap / N / Y
<u>Manuel Baldinu</u> et al [8]	Strangulated right inguinal hernia	Intraoperative / Pyogenic inflammation infiltrated the posterior spermatic cord (Type 3)	Open / N / Y
Tsalis K et al [9]	Recurrent right inguinal hernia	Pre-operative (CECT) / non inflamed (Type 1)	Lap / Y / N
<u>Atsushi Okita</u> [10]	Right incarcerated Indirect inguinal hernia	Intraoperative / Non inflamed (Type 1)	Open / Y / Y
<u>Jayateertha Joshi</u> et al [11]	Left irreducible indirect inguinal hernia	Intraoperative / Non inflamed (Type 1)	Open / N / N
<u>Khalid Alyahyaw</u> [12]	Incarcerated left sided inguinal hernia	Intraoperative / Non inflamed (Type 1)	Open / Y / N
<u>Faranak Olamaeian</u> et al [13]	Right sided inguinal hernia	Intraoperative / Inflamed (Type 2)	Open / Y / Y

Table 2: Named abdominal/groin hernias with unusual contents

Sr. No	Name of Hernia	Content	Named after	Year of Discovery
1	Richter's Hernia	Only part of the bowel wall involved	Hans Richter, Germany	1700s
2	Maydl's Hernia	Loop of bowel protruding through multiple defects	Otto Maydl, Austria	1900
3	Amyand's Hernia	Typically, inguinal hernia where the appendix is the content, can occur in other locations as well	Claudius Amyand, Britain	1735
4	Gibbon's Hernia	Associated with a hydrocele	Edward Gibbon, English historian who suffered from it	1780s
5	Littre Hernia	Inguinal hernia where Meckel's diverticulum is the content	René Litré, France	1700
6	De Garengeot hernia	Femoral hernia where the appendix is the content	René-Jacques De Garengeot, France	1731
7	Narath Hernia	Femoral hernia where hernial sac protrudes behind the femoral vessels in the groin, typically occurring in individuals with congenital hip dislocation	Albert Narath, France	1949
8	Laugier Hernia	Hernial sac pushes through the lacunar ligament into the thigh	Henrie de Laugier, France	1833
9	Serafini Hernia	Femoral hernia in which hernial sac emerges behind femoral vessels	Luigi Serafini, Italy	
10	Velpeau Hernia	Femoral hernia in which hernial sac lies in front of the femoral blood vessels	Alfred Velpeau, France	1820s
11	Cloquets Hernia	Femoral hernia in which hernial sac protrudes through the pectineal fascia, essentially passing over the pectineus muscle and into the thigh	Jules Germain Cloquet, France	1820s
12	Beclard's Hernia	Femoral hernia in which hernial sac protrudes through the saphenous opening	Pierre Augustin Beclard, France	1825
13	Grynfeltt-Lesshaft Hernia	Occurs in the superior lumbar triangle (lined by the 12th rib, the internal oblique muscle, and the erector spinae muscle)	Franz Grynfeltt & Johann Friedrich Lesshaft Germany	1854
14	Petit Hernia	Occurs in the inferior lumbar triangle (lined by the iliac crest, external oblique muscle, and latissimus dorsi muscle)	Jean Louis Petit, France	1743
15	Spigelian Hernia	Occurs through the Spigelian fascia, which is a layer of connective tissue located between the muscles of the abdomen, specifically in the area of the semilunar line	Adriaan van den Spieghel, Holland	1627

CONCLUSION

Amyand's hernia, though rare, should be considered during hernia repair, especially when unusual intraoperative findings are encountered. Laparoscopic approaches allow for effective diagnosis and tailored treatment. In cases where the appendix is normal, reduction and mesh repair yield excellent outcomes.

REFERENCES

- Kenana Tawashi, Eva Khalouf, Muhannad Debes Wzet, *et al.* Intraoperative diagnosis of Amyand's hernia, a case report, *Int. J. Surg. Case Rep.* 2024; 124. doi:202410.1016/j.ijscr.2024.110465.
- Ivanschuk G, Cesmebasi A, Sorenson EP, *et al.* Amyand's hernia: a review. *Med Sci Monit.* 2014; 20:140-6. doi: 10.12659/MSM.889873.
- Kasra Hatampour, Amir Zamani, Roozbeh Shadidi Asil, *et al.* Amyand's hernia in an elective inguinal hernia repair: A case report, *Int. J. Surg. Case Rep.* 2023; 110. doi: 10.1016/j.ijscr.2023.108699.
- Georgia Fezoulidi, Vasiliki Argyrouli, Evangelos Adamopoulos, *et al.* Amyand's hernia: presumptive diagnosis by CT and literature review. *Rad Case Rep.* 2021; 16(4):911-915. doi: 10.1016/j.radcr.2021.01.057.
- Singal R, Gupta S. "Amyand's Hernia" - Pathophysiology, Role of Investigations and Treatment. *Maedica (Bucur).* 2011; 6(4):321-7.
- Khalid H, Khan NA, Aziz MA. Amyand's hernia a case report. *Int J Surg Case Rep.* 2021; 86:106332. doi: 10.1016/j.ijscr.2021.106332.
- Wu SY, Wu ST. Amyand's Hernia. *Asian J Surg.* 2024; S1015-9584(24)0181:8-9. doi: 10.1016/j.asjsur.2024.08.077.
- Montali F, Baldinu M, Annicchiarico A, *et al.* Amyand's hernia: which oncologic risk can be hidden in the sac? *Acta Biomed.* 2022; 93(S1):e2022273. doi: 10.23750/abm.v93iS1.13322.
- Tsalis K, Symeonidis S, Anestiadou E, *et al.* Amyand's Hernia as a Random Finding in Acute Abdominal Pain and the Role of Thorough Investigation: a Case Report. *Maedica (Bucur).* 2022; 17(3):720-725. doi: 10.26574/maedica.2022.17.3.720.
- Okita A, Yoshida O, Murakami M. Incarcerated Amyand's Hernia. *Acta Med Okayama.* 2020; 74(2):171-174. doi: 10.18926/AMO/58277.
- Joshi J, Mallik E, Ahmed T, *et al.* Left sided Amyand hernia - A case report. *Int J Surg Case Rep.* 2022; 96:107374. doi: 10.1016/j.ijscr.2022.107374.
- Alyahyawi K. Left-Sided Amyand's Hernia: A Rare Variant of Inguinal Hernia. *Cureus* 2023; 15(9): e45113. doi: 10.7759/cureus.45113
- Olamaeian F, Saberi Pirouz M, Sheibani F, *et al.* Amyand's hernia: non incarcerated, inflamed appendix in inguinal sac case report. *J Surg Case Rep.* 2022; 2022(9):382. doi: 10.1093/jscr/rjac382.

Funding: Nil; Conflicts of Interest: None Stated.

How to cite this article: Agrawal PP, Joshi AS. Laparoscopic repair of Amyand hernia – A case report with review of literature. *Eastern J Med Sci.* 2025; 10(3):106-109.

