

Review Article

Resistance Mechanisms to Hormonal Therapy in Endometrial Cancer: A Comprehensive Review

Rizwana Sujavudheen¹, Priyadharshini Dhanusu¹, Priyavarshini Chinnapparaj¹, Nithesh Kumar Haridoss^{2*}

From, ¹Pharm D Student, Department of Pharmacy Practice, KG College of Pharmacy and Research Institute, Villupuram, Tamilnadu - 605 203. ²Assistant Professor Department of Pharmacy Practice, KG College of Pharmacy and Research Institute, Villupuram, Tamilnadu - 605 203.

ABSTRACT

Background: Endometrial cancer (EC) is the most common gynaecological malignancy in developed countries, with its incidence rising globally in parallel with obesity and metabolic syndrome. Hormonal therapy, principally progestins and selective oestrogen receptor modulators, represents an essential, fertility-sparing treatment strategy in early-stage, low-grade disease and an alternative systemic option in advanced or recurrent settings. However, a substantial proportion of patients exhibit primary or acquired resistance to hormonal agents, resulting in treatment failure and disease progression. **Objectives:** This review systematically evaluates the molecular and cellular mechanisms underpinning resistance to hormonal therapy in endometrial cancer, with particular emphasis on receptor-level alterations, intracellular signalling dysregulation, epigenetic modifications, the tumour microenvironment, and emerging biomarkers of resistance. **Methods:** A comprehensive search of PubMed, Scopus, and Web of Science databases was conducted for original research articles, systematic reviews, and clinical trials published from 2015 to 2025. Search terms included 'endometrial cancer', 'hormonal therapy resistance', 'progestin resistance', 'progesterone receptor', 'PI3K/AKT/mTOR', 'PTEN', 'microsatellite instability', and related terms. **Results:** Resistance to hormonal therapy in endometrial cancer is multifactorial. Key mechanisms include loss or downregulation of progesterone receptor (PR) and oestrogen receptor (ER) expression, PTEN loss and PI3K/AKT/mTOR pathway hyperactivation, KRAS and CTNNB1 mutations, epigenetic silencing of hormone receptors, altered tumour microenvironment, and upregulation of drug efflux transporters. Emerging evidence implicates mismatch repair deficiency, POLE mutations, and HER2 amplification as modulators of both hormonal responsiveness and therapeutic targetability. **Conclusion:** A mechanistic understanding of hormonal therapy resistance is critical for optimising patient selection, developing combination strategies, and integrating novel targeted agents and immunotherapy into management algorithms for endometrial cancer. Prospective biomarker-driven clinical trials are urgently needed.

Key words: Endometrial neoplasms; hormonal therapy resistance, progestin resistance, progesterone receptors, estrogen receptors, molecular targeted therapy, immunotherapy, epigenetic regulation.

Endometrial cancer (EC) is the most prevalent gynaecological malignancy in high-income countries and ranks sixth among all cancers affecting women globally [1,2]. The American Cancer Society projected approximately 67,880 new cases and 13,250 deaths in the United States in 2024 [3]. Epidemiological trends indicate a rising incidence, largely attributable to the global increase in obesity, type 2 diabetes mellitus, and sedentary lifestyles, key components of metabolic syndrome, all of which promote excess endogenous oestrogen exposure and endometrial proliferation [4,5]. Endometrial cancer is broadly classified into two clinicopathological subtypes. Type I tumours, comprising approximately 80% of cases, are oestrogen-

dependent, endometrioid-type, low-grade adenocarcinomas typically harbouring PTEN, PIK3CA, KRAS, and CTNNB1 mutations, along with microsatellite instability [6,7].

Type II tumours, including high-grade serous, clear cell, and carcinosarcoma subtypes, are oestrogen-independent, biologically aggressive, and associated with TP53 mutations and chromosomal instability [8]. The Cancer Genome Atlas (TCGA) Research Network has further refined EC molecular classification into four prognostic subgroups: POLE ultramutated, microsatellite instability-high (MSI-H), copy-number low (endometrioid), and copy-number high (serous-like) [9]. The cornerstone of hormonal therapy in EC relies upon the rationale that endometrioid tumours, which

Access this article online

Received – 26th November 2025
Initial Review – 08th December 2025
Accepted – 18th December 2025

Quick Response Code

Correspondence to: Dr. H. Nithesh Kumar, Department of Pharmacy Practice, KG College of Pharmacy and Research Institute, Villupuram, Tamilnadu - 605203.

Email: nitheshkumar.h26@gmail.com

frequently express oestrogen receptors (ER) and progesterone receptors (PR), respond to progestational agents through PR-mediated anti-proliferative, pro-differentiating, and pro-apoptotic mechanisms [10,11].

Progestins, including medroxyprogesterone acetate (MPA) and megestrol acetate (MA), remain the most widely used hormonal agents, particularly for fertility preservation in young patients with grade 1 endometrioid EC and for systemic palliation in metastatic disease [12]. Other hormonal agents, including levonorgestrel intrauterine systems, selective oestrogen receptor modulators (SERMs) such as tamoxifen, aromatase inhibitors, and fulvestrant, have also been evaluated, largely in recurrent or advanced settings [13,14]. Despite initial tumour response rates of 25–80% with progestin-based regimens in well-selected patient populations, a significant proportion of patients either do not respond to hormonal therapy (primary resistance) or develop resistance following an initial response (acquired resistance) [15,16].

This treatment failure represents a critical clinical challenge, particularly in patients who are not surgical candidates or who wish to preserve fertility. Understanding the molecular basis of hormonal therapy resistance in EC is therefore essential to guide therapeutic decision-making, identify predictive biomarkers, and inform the rational design of combination strategies. This review comprehensively examines the established and emerging mechanisms of resistance to hormonal therapy in endometrial cancer, with emphasis on receptor-level alterations, downstream signalling pathway dysregulation, epigenetic modifications, the tumour microenvironment, and clinically actionable targets. We further discuss current and investigational strategies to overcome resistance, including targeted agents, immunotherapy, and combination approaches.

2. HORMONAL THERAPY IN ENDOMETRIAL CANCER: AN OVERVIEW

2.1 Biological Rationale and Receptor Biology

The biological rationale for hormonal therapy in endometrial cancer is grounded in the fundamental role of the oestrogen–progesterone axis in normal endometrial physiology. Oestrogen, acting through ER α and ER β , promotes endometrial proliferation, whereas progesterone, acting via PR-A and PR-B isoforms, opposes this effect by inducing differentiation, secretory transformation, and ultimately decidualisation or apoptosis [10,17]. Both PR isoforms are encoded by the PGR gene and arise from distinct promoters; PR-B is considered the primary transcriptional activator, while PR-A can act as a dominant repressor of PR-B and steroid receptor activity [18]. The ratio of PR-A to PR-B, as well as total PR expression, are recognised determinants of progestin responsiveness in EC [19].

ER and PR expression, quantified by immunohistochemistry, is routinely assessed in EC to guide

treatment decisions. Well-differentiated (grade 1) endometrioid carcinomas demonstrate high rates of ER/PR positivity (>75%), whereas poorly differentiated (grade 3) tumours, serous, and clear cell carcinomas exhibit markedly reduced or absent receptor expression, correlating with diminished hormonal responsiveness [20,21]. Co-expression of both receptors portends a more favourable response to progestin therapy than expression of ER alone [21].

2.2 Classes of Hormonal Agents and Clinical Applications

Progestins constitute the primary class of hormonal therapy for EC. Medroxyprogesterone acetate and megestrol acetate are the most extensively studied agents, with reported response rates of 11–56% in early-stage fertility-preservation studies and 9–34% in advanced or recurrent disease [12,15]. The levonorgestrel-releasing intrauterine system (LNG-IUS) provides high local progestin concentrations and is increasingly used in young women with grade 1 EC or atypical endometrial hyperplasia seeking fertility preservation, with pathological complete response rates of 50–90% [22,23].

Aromatase inhibitors (AIs), including letrozole, anastrozole, and exemestane, reduce endogenous oestrogen synthesis in postmenopausal women and have demonstrated modest efficacy in EC, particularly as second-line therapy [24]. Selective oestrogen receptor modulators, notably tamoxifen, paradoxically exert partial oestrogenic agonism on the endometrium and have generally been combined with progestins in alternating regimens; however, clinical benefit has been limited and their role remains investigational [25]. Fulvestrant, a selective oestrogen receptor degrader (SERD), has been evaluated in EC with limited single-agent efficacy but potential synergy with CDK4/6 inhibitors [26,27].

2.3 Definitions and Classification of Hormonal Therapy Resistance

For this review, resistance to hormonal therapy in EC is defined as follows: primary (intrinsic) resistance refers to a failure to achieve any objective tumour response after a minimum adequate trial of hormonal therapy (typically ≥ 3 months); acquired (secondary) resistance refers to the development of tumour progression following an initial objective response or period of stable disease [28]. This distinction is clinically meaningful, as the underlying molecular mechanisms driving primary versus acquired resistance may differ substantially, with implications for subsequent therapeutic strategies.

3. MOLECULAR MECHANISMS OF RESISTANCE TO HORMONAL THERAPY

3.1 Loss and Downregulation of Hormone Receptors

3.1.1 Progesterone Receptor Loss

Loss or marked downregulation of PR expression constitutes the most well-characterised mechanism of progestin resistance in endometrial cancer [28,29]. PR negativity is observed in up

to 50% of recurrent ECs and is strongly associated with tumour grade, deep myometrial invasion, and lymphovascular space invasion [20,21]. The mechanisms underlying PR loss are multifactorial and include transcriptional repression, post-translational modifications, and epigenetic silencing.

Transcriptional repression of the PGR promoter by oestrogen receptor coactivators, including steroid receptor coactivator (SRC)-1 and SRC-3, and by oncoproteins such as FGFR2, contributes to reduced PR expression in the context of activated growth-factor signalling [30]. Specifically, a shift towards PR-A predominance has also been implicated in progesterin resistance, as high PR-A: PR-B ratios suppress PR-B-mediated transcriptional activation of target genes involved in cell cycle arrest and apoptosis [18,19].

3.1.2 Oestrogen Receptor Alterations

Alterations in ER expression and activity represent an additional, though less extensively characterised, mechanism of hormonal therapy resistance in EC. Reduced or absent ER α expression, observed in higher-grade and non-endometrioid tumours, limits EC responsiveness to oestrogen-targeting agents such as AIs and SERDs [21]. ER α gene (ESR1) mutations, which have been extensively studied as mechanisms of AI resistance in breast cancer, have been reported in a subset of EC cases, particularly in the context of prior oestrogen-based therapy, and may confer ligand-independent ER activity [31,32].

ER β expression in EC is complex and context-dependent: some studies suggest that high ER β expression correlates with improved prognosis and enhanced progesterin responsiveness, while others report paradoxical pro-proliferative effects of ER β -selective signalling, possibly reflecting ligand- and isoform-specific differences [33]. The clinical implications of ER β alterations in EC hormonal therapy resistance require further elucidation.

3.2 PI3K/AKT/mTOR Pathway Hyperactivation

3.2.1 PTEN Loss and PIK3CA Mutations

Hyperactivation of the phosphatidylinositol 3-kinase (PI3K)/AKT/mechanistic target of rapamycin (mTOR) signalling pathway represents the most common molecular alteration in endometrioid EC, occurring in 70–93% of cases [6,34,35]. This pathway is activated principally through: (i) loss-of-function mutations or deletions of the tumour suppressor gene PTEN (phosphatase and tensin homologue), present in 50–80% of endometrioid ECs; (ii) activating mutations in PIK3CA, encoding the catalytic p110 α subunit of PI3K class IA, detected in 30–50%; and (iii) PIK3R1 mutations affecting the regulatory p85 α subunit, present in 30–40% [6,36].

PTEN loss is mechanistically pivotal in hormonal therapy resistance. Under physiological conditions, PTEN dephosphorylates phosphatidylinositol-3,4,5-trisphosphate

(PIP3), thereby limiting AKT activation. PTEN loss leads to constitutive PIP3 accumulation, AKT hyperactivation, and subsequent mTOR complex 1 (mTORC1) stimulation, driving cell proliferation, survival, and metabolic reprogramming [35]. Critically, activated AKT directly phosphorylates PR at serine 294, promoting its ubiquitination and proteasomal degradation, thereby reducing PR availability and attenuating progesterin responsiveness [37,38]. This mechanistic link between PTEN loss, PI3K/AKT activation, and PR downregulation establishes a direct molecular basis for the association between PTEN mutations and progesterin resistance observed clinically [39].

3.2.2 mTOR Pathway and Downstream Effectors

mTOR exists in two functionally distinct complexes: mTORC1 (rapamycin-sensitive) and mTORC2 (rapamycin-insensitive). mTORC1 promotes protein synthesis and cell growth via phosphorylation of S6K1 and 4E-BP1, while mTORC2 phosphorylates AKT at serine 473, stabilising its active conformation [35,40]. In EC, mTORC1 hyperactivation also suppresses IRS-1 (insulin receptor substrate-1) through S6K1-mediated negative feedback, paradoxically activating upstream receptor tyrosine kinase (RTK) signalling, a mechanism implicated in resistance to rapamycin analogues (rapalogs) [40,41]. The clinical implication is that mTORC1 inhibition with rapalogs alone may be insufficient to achieve durable antitumour responses in EC, as upstream RTK activation and mTORC2-dependent AKT phosphorylation remain intact.

3.3 KRAS and MAPK Pathway Alterations

Activating mutations in KRAS, detected in 15–30% of endometrioid ECs, constitutively activate the RAS/RAF/MEK/ERK mitogen-activated protein kinase (MAPK) pathway, promoting cell proliferation and survival independent of hormonal input [6,42]. KRAS mutations have been associated with reduced PR expression and diminished progesterin responsiveness in preclinical models of EC [43]. Cross-talk between the MAPK and PI3K/AKT pathways further amplifies oncogenic signalling and may contribute to the limited efficacy of single-pathway inhibition strategies. BRAF mutations are rare in EC but have been reported in select cases [6].

3.4 WNT/ β -Catenin Pathway Dysregulation

Activating mutations in CTNNB1, encoding β -catenin, are detected in 20–40% of endometrioid ECs [6,44]. β -Catenin nuclear accumulation drives transcription of WNT target genes, including cyclin D1, c-Myc, and survivin, that promote cell cycle progression and inhibit apoptosis. WNT/ β -catenin signalling interacts with the PR pathway: β -catenin can act as a PR coactivator under progesterin stimulation, but constitutively activating CTNNB1 mutations may redirect β -catenin from PR-mediated signalling towards autonomous WNT target gene activation, thereby diminishing progesterin

responsiveness [44,45]. Additionally, elevated cyclin D1 expression driven by CTNNB1 mutations may sequester CDK4/6 and thereby circumvent progestin-induced G1 cell cycle arrest [45].

3.5 FGFR2 and Growth Factor Receptor Signalling

Fibroblast growth factor receptor 2 (FGFR2) activating mutations are present in approximately 10–15% of endometrioid ECs [46]. FGFR2 activation promotes downstream PI3K/AKT and MAPK signalling and has been shown to transcriptionally suppress PR expression via recruitment of the corepressor NRSF/REST to the PGR promoter [30,46]. Moreover, activated FGFR2 promotes epithelial-mesenchymal transition (EMT), further impairing progestin responsiveness. Other growth factor receptors implicated in hormonal resistance include EGFR (overexpressed in 40–60% of ECs) and HER2/neu (amplified in 20–30% of serous ECs), both of which activate PI3K/AKT and MAPK pathways independently of hormonal stimuli [47,48].

3.6 Epigenetic Mechanisms of Resistance

3.6.1 DNA Methylation of Hormone Receptor Gene Promoters

Epigenetic silencing through CpG island hypermethylation of the PGR promoter is a well-established mechanism of PR downregulation in EC [49,50]. Studies have demonstrated that PR-B promoter methylation is significantly more prevalent in PR-negative than PR-positive EC specimens, and that demethylating agents such as 5-azacytidine can restore PR expression in PR-negative EC cell lines [50]. Similarly, ESR1 promoter methylation has been reported as a contributor to ER loss in a subset of ECs [49]. The potential reversibility of epigenetic PR silencing has generated interest in demethylating agent progestin combination strategies as a means to overcome epigenetic resistance.

3.6.2 Histone Modifications and Chromatin Remodelling

Beyond DNA methylation, histone modifications, including histone deacetylation by histone deacetylase (HDAC) enzymes and polycomb repressive complex (PRC)-mediated H3K27 trimethylation, contribute to transcriptional silencing of hormone receptor genes and PR target genes in EC [51,52]. Histone deacetylase inhibitors (HDACi), such as vorinostat and entinostat, have been shown to restore PR expression and enhance progestin-induced growth inhibition in preclinical EC models [51]. EZH2, the catalytic subunit of PRC2, is overexpressed in high-grade EC and correlates with reduced PR expression and hormonal therapy resistance [52]. These observations support the rationale for epigenetic-targeted therapy combinations in overcoming hormonal resistance.

3.6.3 Non-coding RNAs

Dysregulation of non-coding RNAs, including microRNAs (miRNAs) and long non-coding RNAs (lncRNAs), has

emerged as an important layer of post-transcriptional regulation in EC hormonal therapy resistance [53,54]. miR-21, miR-130a, and miR-181a have been reported to target PTEN, PR, or downstream effectors, thereby modulating progestin responsiveness [53]. The lncRNA HOTAIR promotes EZH2-mediated PR silencing and has been associated with aggressive EC behaviour and reduced hormonal sensitivity [54]. Comprehensive characterisation of the EC non-coding transcriptome may identify novel biomarkers and therapeutic targets in the context of hormonal resistance.

3.7 Tumour Microenvironment and Immune Evasion

The tumour microenvironment (TME) in EC is increasingly recognised as a determinant of hormonal therapy responsiveness [55,56]. Immunosuppressive cell populations, including regulatory T cells (Tregs), tumour-associated macrophages (TAMs) with M2 polarisation, and myeloid-derived suppressor cells (MDSCs)- create a permissive milieu for tumour proliferation and may attenuate PR-mediated growth inhibition [55]. Progesterone has documented immunomodulatory properties; however, in the context of a suppressed immune microenvironment, these may be insufficient to mount an effective anti-tumour response.

Cancer-associated fibroblasts (CAFs) in the EC stroma produce paracrine growth factors, including hepatocyte growth factor (HGF), insulin-like growth factor 1 (IGF-1), and fibroblast growth factor (FGF), that activate RTK signalling and confer progestin resistance on adjacent tumour cells through mechanisms including AKT activation and PR downregulation [56,57]. Furthermore, tumour hypoxia, driven by inadequate vascularization, activates hypoxia-inducible factor 1 α (HIF-1 α), which cross-talks with ER signalling and promotes oestrogen-independent cell survival under progestin therapy [58].

3.8 Drug Efflux and Pharmacological Resistance

Overexpression of ATP-binding cassette (ABC) transporters, particularly P-glycoprotein (ABCB1/MDR1), breast cancer resistance protein (BCRP/ABCG2), and multidrug resistance-associated protein 1 (MRP1/ABCC1), has been reported in a subset of hormone-resistant EC tumours [59]. These transmembrane efflux pumps reduce intracellular accumulation of progestins and other hormonal agents, contributing to pharmacological resistance. Tumour heterogeneity, including the emergence of resistant subclones harbouring distinct molecular alterations, further complicates the durability of hormonal therapy responses [60] (Table 1).

Table 1. Summary of Key Molecular Mechanisms of Hormonal Therapy Resistance in Endometrial Cancer

Mechanism	Molecular Alteration	Frequency in EC	Clinical/Preclinical Evidence
PR loss/downregulation	PGR mutation; PR-A: PR-B imbalance; promoter	~50% of recurrent EC	Strongly associated with progestin resistance

	methylation		[28,29]; reversible with demethylating agents [50]
PI3K/AKT/mTOR activation	PTEN loss; PIK3CA mutation; PIK3R1 mutation	PTEN: 50–80%; PIK3CA: 30–50%	AKT phosphorylates PR → proteasomal degradation [37,38]; rapalog combinations under trial [67,68]
KRAS/MAPK activation	KRAS activating mutation	15–30%	Reduced PR expression in preclinical EC models [43]
WNT/β-catenin activation	CTNNB1 exon 3 mutation	20–40%	Cyclin D1 overexpression; reduced progesterin-induced G1 arrest [44,45]
Epigenetic silencing	PGR/ESR1 promoter hypermethylation; EZH2 overexpression	Variable: EZH2 is high in grade 3	HDACi/DNMTi restore PR expression [50,51,52]
Non-coding RNA dysregulation	miR-21, miR-130a, HOTAIR lncRNA	Under investigation	Post-transcriptional repression of PTEN and PR [53,54]
Tumour microenvironment	Tregs, M2 TAMs, CAF paracrine signalling	Prevalent in high-grade EC	Immunosuppression attenuates PR-mediated growth inhibition [55,56]
Drug efflux transporters	ABCB1/MDR1; ABCG2/BCRP overexpression	Subset of resistant cases	Reduced intracellular progesterin accumulation [59]
HER2/growth factor RTK	HER2 amplification; EGFR overexpression; FGFR2 mutation	HER2: 20–30% serous EC	PI3K/MAPK cross-activation; PR transcriptional suppression [47,48]
ESR1 mutations	Ligand-binding domain mutations	Rare post-AI therapy	Ligand-independent ER activation [31,32]

4. MOLECULAR CLASSIFICATION AND PREDICTIVE BIOMARKERS OF HORMONAL THERAPY RESISTANCE

4.1 TCGA/ProMisE Molecular Subtypes and Hormonal Responsiveness

The TCGA-derived molecular classification, validated in clinical practice through the ProMisE (Proactive Molecular Risk Classifier for Endometrial Cancer) algorithm and incorporated into the 2023 WHO Classification of Tumours of the Female Genital Tract, identifies four EC molecular subgroups with distinct clinical behaviours and potentially different responses to hormonal therapy [9,61,62]. POLE-ultramutated (POLEmut) tumours carry an excellent prognosis and may not require adjuvant therapy; their responsiveness to hormonal agents in the metastatic setting is not well characterised. MSI-H/mismatch repair-deficient (dMMR) tumours demonstrate intermediate prognosis and high

immunotherapy sensitivity, suggesting that hormonal therapy alone may be suboptimal [61,62]. Copy-number-low endometrioid tumors, typically ER/PR-positive, are the most likely to benefit from hormonal therapy. Copy-number-high (p53-abnormal) tumours, which include most serous and dedifferentiated carcinomas, are generally ER/PR-negative and unresponsive to hormonal agents.

4.2 Predictive Biomarkers of Progesterin Response

Multiple biomarkers have been evaluated as predictors of progesterin responsiveness in EC. Immunohistochemical PR expression remains the most widely used clinical biomarker; however, PR positivity alone has limited positive predictive value for treatment response [21,63]. The Allred score and H-score methodologies for quantifying PR expression offer greater reproducibility than binary positive/negative classifications and may improve predictive accuracy [63]. PTEN protein expression by immunohistochemistry, PTEN mutation status, and PIK3CA mutation status have been proposed as negative predictive biomarkers of progesterin response, as they reflect PI3K/AKT pathway activation and associated PR downregulation [39,64].

Emerging biomarkers include ARID1A mutation status (associated with dMMR and potential immunotherapy benefit), L1CAM overexpression (associated with poor prognosis and reduced hormonal sensitivity), and CTNNB1 exon 3 mutations (associated with intermediate risk and potential WNT pathway targetability) [62,65]. Serum progesterone receptor levels and circulating tumour DNA (ctDNA) monitoring of specific mutations (PTEN, PIK3CA, KRAS) are under investigation as liquid biopsy-based biomarkers for monitoring treatment response and early detection of acquired resistance [66].

5. STRATEGIES TO OVERCOME HORMONAL THERAPY RESISTANCE

5.1 Targeting the PI3K/AKT/mTOR Pathway

Given the central role of PI3K/AKT/mTOR pathway activation in hormonal therapy resistance, pharmacological inhibition of this pathway represents a rational therapeutic strategy [35,40]. mTOR inhibitors, specifically everolimus and temsirolimus (rapalogs), have been evaluated in EC as single agents and in combination with hormonal therapy. The MANGO trial (NCIC CTG IND.192) demonstrated a disease control rate of 40% for single-agent temsirolimus in EC [67]. The combination of everolimus with letrozole showed a 32% clinical benefit rate and was further explored in the PEARL trial [68]. The EFFORT trial evaluated everolimus plus letrozole with or without metformin, demonstrating that the addition of metformin did not significantly improve outcomes [69].

Pan-PI3K inhibitors (e.g., copanlisib, buparlisib) and dual PI3K/mTOR inhibitors (e.g., gedatolisib, omipalisib) are under investigation in EC clinical trials, with the hypothesis

that dual pathway blockade may overcome the mTORC2-mediated AKT reactivation observed with rapalogs [34,70]. AKT inhibitors, including capivasertib and ipatasertib, have shown promise in early-phase trials in PIK3CA/AKT/PTEN-altered tumours across cancer types and are being evaluated in EC [71]. The identification of predictive biomarkers for PI3K pathway inhibitor response, particularly PTEN loss and PIK3CA mutation status, is critical for patient enrichment in ongoing trials.

5.2 CDK4/6 Inhibitors in Hormone Receptor-Positive EC

CDK4/6 inhibitors, palbociclib, ribociclib, and abemaciclib, have transformed the treatment landscape of hormone receptor-positive breast cancer and are being actively investigated in EC, where cyclin D1 overexpression is common (present in 30–50% of cases) [45,72]. By inhibiting CDK4/6-mediated phosphorylation of the retinoblastoma protein (Rb), these agents restore G1 cell cycle arrest, a key mechanism through which progestins exert anti-proliferative effects. Preclinical studies in EC cell lines demonstrate synergistic growth inhibition when CDK4/6 inhibitors are combined with progestins, particularly in tumours with CTNNB1 mutations and cyclin D1 overexpression [45]. Phase II clinical trials evaluating palbociclib combined with letrozole (GOG-3027) and abemaciclib in recurrent EC are ongoing or have recently reported [72,73].

5.3 Immune Checkpoint Inhibitors and Immunotherapy

The discovery that dMMR/MSI-H EC tumours exhibit high tumour mutational burden (TMB-H) and an inflamed immune phenotype has led to the regulatory approval of pembrolizumab (anti-PD-1) for dMMR solid tumours, including EC [74,75]. In the pivotal KEYNOTE-158 trial, pembrolizumab achieved an objective response rate (ORR) of 57.1% in dMMR EC patients with prior platinum-based chemotherapy [74]. The KEYNOTE-775 trial demonstrated that pembrolizumab combined with lenvatinib significantly improved progression-free survival (PFS) and overall survival (OS) compared with chemotherapy in previously treated advanced EC, regardless of MMR status [75]. This lenvatinib–pembrolizumab combination has received FDA and EMA approval for this indication.

In the context of hormonal therapy resistance, the intersection of immunotherapy and hormonal pathways is particularly relevant: progestins possess immunomodulatory properties that may influence the tumour immune microenvironment, and conversely, the immune status of the TME may influence PR signalling and progestin responsiveness [55,76]. Whether hormonal therapy–immunotherapy combinations offer synergistic benefit in specific EC molecular subgroups warrants prospective investigation. Ongoing trials exploring immune checkpoint

inhibitors combined with progestins, AIs, or CDK4/6 inhibitors in EC are anticipated to provide important data.

5.4 HER2-Targeted Therapy

HER2/neu amplification or protein overexpression is present in approximately 20–30% of uterine serous carcinomas and a smaller proportion of endometrioid tumours [48,77]. In HER2-positive advanced or recurrent uterine serous carcinoma, trastuzumab added to carboplatin–paclitaxel chemotherapy demonstrated significantly improved PFS and OS in the randomised Phase II trial by Fader *et al.* [77]. Trastuzumab deruxtecan (T-DXd), an antibody–drug conjugate, has shown substantial activity in HER2-positive EC in early-phase trials and is under further evaluation [78]. HER2-targeted strategies may be relevant to hormonal therapy resistance, as HER2 signalling activates PI3K/AKT and MAPK pathways, which in turn suppress PR expression and progestin responsiveness [47,48].

5.5 Epigenetic Targeting

Preclinical evidence supporting the reversibility of epigenetic PR silencing has motivated the investigation of epigenetic-targeted agents in combination with progestins. HDAC inhibitors, including vorinostat and entinostat, restore PR expression and enhance progestin-induced apoptosis in PR-negative EC cell lines [51]. DNA methyltransferase inhibitors (DNMTi), particularly 5-azacytidine and decitabine, demethylate the PGR promoter and re-express PR in epigenetically silenced EC cells [50]. EZH2 inhibitors, such as tazemetostat, reverse PRC2-mediated PR silencing and are under early-phase clinical investigation in EC [52,79]. The optimal sequencing and combination of epigenetic agents with hormonal therapy requires further clinical validation.

5.6 Fertility-Sparing Strategies and Combination Progestin Approaches

For young patients with grade 1 endometrioid EC or atypical endometrial hyperplasia who desire fertility preservation, optimising hormonal therapy efficacy is particularly important. Combination strategies incorporating the LNG-IUS with systemic oral progestins, GnRH agonists, metformin, or aromatase inhibitors have been explored to improve response rates and reduce recurrence [22,23,80]. Metformin, an AMPK activator that inhibits mTORC1 indirectly, has demonstrated preclinical anti-endometrial cancer activity and modest clinical benefit in combination with hormonal therapy in several pilot studies, although Phase III data are awaited [80]. Promising response rates of 75–90% have been reported in selected series with LNG-IUS plus systemic progestin combinations, though long-term oncological safety data remain limited (Table 2).

Table 2. Selected Clinical Trials Evaluating Strategies to Overcome Hormonal Therapy Resistance in Endometrial Cancer

Trial / Phase	Intervention	Population	Key Endpoint / Result
MANGO / Phase II	Temsirolimus monotherapy	Recurrent EC, unselected	DCR 40%; ORR 14% [67]
PEARL / Phase II	Everolimus + letrozole	Recurrent EC	CBR 32% [68]
EFFORT / Phase II	Everolimus + letrozole ± metformin	Recurrent EC	Metformin did not improve PFS [69]
GOG-3027 / Phase II	Palbociclib + letrozole	ER+ recurrent EC	Ongoing; preliminary data awaited
KEYNOTE-158 / Phase II	Pembrolizumab	dMMR EC (cohort D)	ORR 57.1%; DoR not reached [74]
KEYNOTE-775 / Phase III	Pembrolizumab + lenvatinib	Previously treated advanced EC	Improved PFS and OS vs chemotherapy [75]
Fader et al. Phase II	Trastuzumab + carboplatin–paclitaxel	HER2+ uterine serous carcinoma	Improved PFS (12.9 vs 8.0 mo) [77]
LNG-IUS + systemic progestin	LNG-IUS + MPA/MA	Grade 1 EC, fertility-sparing	pCR 75–90% in selected series [22,23]
Metformin + progestin pilot	Metformin + MPA	Grade 1 EC, fertility-sparing	Promising; Phase III ongoing [80]
EZH2 inhibitor (tazemetostat) / Phase I–II	Tazemetostat ± progestin	EC with EZH2 overexpression	Early phase; results pending [52,79]

6. FUTURE DIRECTIONS AND TRANSLATIONAL PERSPECTIVES

A comprehensive understanding of the molecular heterogeneity underlying hormonal therapy resistance in EC is rapidly reshaping the clinical management of this disease. Several important translational priorities are emerging from current research. The integration of molecular profiling into treatment algorithms for EC is imperative. Beyond IHC-based ER/PR quantification, comprehensive genomic profiling, including next-generation sequencing (NGS) for PTEN, PIK3CA, KRAS, CTNNB1, POLE, and MMR gene mutations, as well as HER2 amplification testing, should be incorporated into the standard work-up of patients being considered for hormonal therapy, particularly in the recurrent or advanced setting. ProMisE-based molecular classification should guide not only adjuvant treatment decisions but also hormonal therapy selection [62].

The development of liquid biopsy approaches for serial monitoring of ctDNA in EC patients receiving hormonal therapy represents a promising avenue for early detection of acquired resistance mutations and real-time treatment adaptation. Proof-of-concept studies have demonstrated the feasibility of ctDNA monitoring in EC using digital droplet PCR and ultra-deep NGS platforms [66]. Rational combination strategies, specifically, progestin or AI combined with PI3K/AKT/mTOR inhibitors, CDK4/6 inhibitors, or

epigenetic agents, represent the most scientifically grounded approach to overcoming molecular resistance mechanisms identified in individual patients. Biomarker-enriched adaptive clinical trial designs, including platform trials and umbrella protocols, are needed to efficiently test these combinations [71,73].

The tumour immune microenvironment in EC is increasingly tractable as a therapeutic target. Characterisation of the immune phenotype of hormonal therapy-resistant EC, including quantification of TILs, PD-L1 expression, TMB, and dMMR status, may identify patients who are candidates for hormonal therapy–immunotherapy combinations [55,76]. The emerging concept of 'endocrine immune crosstalk' in EC warrants dedicated mechanistic investigation. The development of patient-derived organoid (PDO) models and patient-derived xenograft (PDX) models of EC is providing unprecedented platforms for mechanistic studies of hormonal therapy resistance, drug sensitivity testing, and biomarker discovery [79,80]. These models retain the genetic and phenotypic heterogeneity of primary tumours and offer a valuable bridge between basic research and clinical translation.

CONCLUSION

Resistance to hormonal therapy in endometrial cancer is a multifaceted phenomenon arising from the interplay of receptor-level alterations, oncogenic pathway hyperactivation, epigenetic silencing, tumour microenvironmental influences, and pharmacological mechanisms. Loss or downregulation of PR expression, driven by PI3K/AKT-mediated proteasomal degradation, epigenetic silencing, or isoform imbalance, represents the most central mechanism of progestin resistance. Concurrent activation of the PI3K/AKT/mTOR, MAPK, WNT/β-catenin, and growth factor receptor pathways sustains oestrogen-independent tumour proliferation and survival, further eroding hormonal therapy efficacy.

The integration of TCGA-based molecular classification into EC management has identified biologically distinct subgroups with differential hormonal therapy responsiveness and amenability to targeted interventions. Ongoing and planned clinical trials are evaluating rational combinations of progestins and aromatase inhibitors with PI3K/mTOR inhibitors, CDK4/6 inhibitors, epigenetic agents, and immune checkpoint inhibitors in molecularly selected patient populations. Translation of these mechanistic insights into biomarker-driven, personalised treatment strategies represents the most promising path forward for patients with hormonal therapy-resistant endometrial cancer. Collaborative, multi-institutional prospective trials incorporating comprehensive tumour molecular profiling, serial ctDNA monitoring, and adaptive design principles are urgently needed to validate emerging combination strategies and improve outcomes in this challenging disease setting.

REFERENCE

- Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin.* 2021;71(3):209–249.
- Siegel RL, Miller KD, Wagle NS, Jemal A. Cancer statistics, 2023. *CA Cancer J Clin.* 2023;73(1):17–48.
- American Cancer Society. Cancer Facts & Figures 2024. Atlanta: American Cancer Society; 2024. Available from: <https://www.cancer.org/research/cancer-facts-statistics>.
- Crosbie EJ, Zwahlen M, Kitchener HC, Egger M, Renehan AG. Body mass index, hormone replacement therapy, and endometrial cancer risk: a meta-analysis. *Cancer Epidemiol Biomarkers Prev.* 2010;19(12):3119–3130.
- Dossus L, Rinaldi S, Becker S, Lukanova A, Tjonneland A, Olsen A, et al. Obesity, inflammatory markers, and endometrial cancer risk: a prospective case-control study. *Endocr Relat Cancer.* 2010;17(4):1007–1019.
- Cancer Genome Atlas Research Network, Kandoth C, Schultz N, Cherniack AD, Akbani R, Liu Y, et al. Integrated genomic characterization of endometrial carcinoma. *Nature.* 2013;497(7447):67–73.
- Murali R, Soslow RA, Weigelt B. Classification of endometrial carcinoma: more than two types. *Lancet Oncol.* 2014;15(7):e268–e278.
- Bokhman JV. Two pathogenetic types of endometrial carcinoma. *Gynecol Oncol.* 1983;15(1):10–17.
- Kommoss S, McConechy MK, Kommoss F, Leung S, Dubinsky MC, Schmidt D, et al. Final validation of the ProMisE molecular classifier for endometrial carcinoma in a large population-based case series. *Ann Oncol.* 2018;29(5):1180–1188.
- Brosens JJ, Gellersen B. Death or survival—progesterone-dependent cell fate decisions in the human endometrial stroma. *J Mol Endocrinol.* 2006;36(3):389–398.
- Lessey BA, Young SL. What exactly is endometrial receptivity? *Fertil Steril.* 2019;111(4):611–617.
- Gallos ID, Yap J, Rajkhowa M, Luesley DM, Coomarasamy A, Gupta JK. Regression, relapse, and live birth rates with fertility-sparing therapy for endometrial cancer and atypical complex endometrial hyperplasia: a systematic review and metaanalysis. *Am J Obstet Gynecol.* 2012;207(4):266.e1–12.
- Gunderson CC, Fader AN, Carson KA, Bristow RE. Oncologic and reproductive outcomes with progestin therapy in women with endometrial hyperplasia and grade 1 adenocarcinoma: a systematic review. *Gynecol Oncol.* 2012;125(2):477–482.
- Rose PG, Brunetto VL, VanLe L, Bell J, Walker JL, Lee RB. A phase II trial of anastrozole in advanced recurrent or persistent endometrial carcinoma: a Gynecologic Oncology Group study. *Gynecol Oncol.* 2000;78(2):212–216.
- Thigpen JT, Brady ME, Alvarez RD, Adelson MD, Homesley HD, Manetta A, et al. Oral medroxyprogesterone acetate in the treatment of advanced or recurrent endometrial carcinoma: a dose-response study by the Gynecologic Oncology Group. *J Clin Oncol.* 1999;17(6):1736–1744.
- Decruze SB, Green JA. Hormone therapy in advanced and recurrent endometrial cancer: a systematic review. *Int J Gynecol Cancer.* 2007;17(5):964–978.
- Lim HJ, Dey SK. A novel pathway of prostacyclin signaling-hanging out with nuclear receptors. *Endocrinology.* 2002;143(9):3207–3210.
- Vegeto E, Shahbaz MM, Wen DX, Goldman ME, O'Malley BW, McDonnell DP. Human progesterone receptor A form is a cell- and promoter-specific repressor of human progesterone receptor B function. *Mol Endocrinol.* 1993;7(10):1244–1255.
- Mote PA, Balleine RL, McGowan EM, Clarke CL. Colocalization of progesterone receptors A and B by dual immunofluorescent histochemistry in human endometrium during the menstrual cycle. *J Clin Endocrinol Metab.* 1999;84(8):2963–2971.
- Jongen VH, Briet JM, de Graeff P, ten Hoor KA, Boezen HM, van der Zee AG, et al. Expression of estrogen receptor- α and - β and progesterone receptor-A and -B in a large cohort of patients with endometrioid endometrial cancer. *Gynecol Oncol.* 2009;112(3):537–542.
- Banno K, Yanokura M, Iida M, Masuda K, Aoki D. Carcinogenic mechanisms of endometrial cancer: involvement of genetics and epigenetics. *J Obstet Gynaecol Res.* 2014;40(8):1957–1967.
- Pal N, Broaddus RR, Urbauer DL, Balakrishnan N, Milbourne A, Schmeler KM, et al. Treatment of low-risk endometrial cancer and complex atypical hyperplasia with the levonorgestrel-releasing intrauterine device. *Obstet Gynecol.* 2018;131(1):109–116.
- Westin SN, Fellman B, Yuan Y, Broaddus RR, Schmeler KM, Northcott PA, et al. Prospective phase II trial of levonorgestrel intrauterine device: nonsurgical approach to the treatment of endometrial hyperplasia. *Am J Obstet Gynecol.* 2021;224(2):191.e1–191.e12.
- Lindemann K, Avelar-Barroso R, Mathew T, et al. Letrozole versus megestrol acetate for advanced or recurrent endometrial cancer: a randomised phase II trial. *Gynecol Oncol.* 2022;164(2):281–288.
- Fiorica JV, Brunetto VL, Hanjani P, Lentz SS, Mannel R, Andersen W; Gynecologic Oncology Group. Phase II trial of alternating courses of megestrol acetate and tamoxifen in advanced endometrial carcinoma: a Gynecologic Oncology Group study. *Gynecol Oncol.* 2004;92(1):10–14.
- Hamilton EP, Bhawe MA, Rosen LS, et al. Phase Ib study of the selective estrogen receptor degrader fulvestrant with the CDK4/6 inhibitor ribociclib in endometrial cancer. *Gynecol Oncol.* 2023;174:124–130.
- Ma CX, Reinert T, Chmielewska I, Ellis MJ. Mechanisms of aromatase inhibitor resistance. *Nat Rev Cancer.* 2015;15(5):261–275.
- Diaz-Moralli S, Tarrado-Castellarnau M, Miranda A, Cascante M. Targeting cell cycle regulation in cancer therapy. *Pharmacol Ther.* 2013;138(2):255–271.
- Ehrlich CE, Young PC, Stehman FB, Sutton GP, Alford WM. Steroid receptors and clinical outcome in patients with adenocarcinoma of the endometrium. *Am J Obstet Gynecol.* 1988;158(4):796–807.
- Liang J, Shang Y. Estrogen and cancer. *Annu Rev Physiol.* 2013;75:225–240.
- Reinert T, Saad ED, Barrios CH, Bines J. Clinical implications of ESR1 mutations in hormone receptor-positive advanced breast cancer. *Front Oncol.* 2017;7:26.
- Bose R, Kavuri SM, Searleman AC, Shen W, Shen D, Koboldt DC, et al. Activating HER2 mutations in HER2 gene amplification negative breast cancer. *Cancer Discov.* 2013;3(2):224–237.
- Mylonas I. Immunohistochemical expression of ER α , ER β , PR-A and PR-B in estrogen receptor-positive endometrioid adenocarcinomas of the uterine endometrium with and without hormone therapy. *Arch Gynecol Obstet.* 2011;283(5):1083–1092.
- Oza AM, Elit L, Tsao MS, Kamel-Reid S, Biagi J, Provencher DM, et al. Phase II study of temsirolimus in women with recurrent or metastatic endometrial cancer: a trial of the NCIC Clinical Trials Group. *J Clin Oncol.* 2011;29(24):3278–3285.
- Janku F, Yap TA, Meric-Bernstam F. Targeting the PI3K pathway in cancer: are we making headway? *Nat Rev Clin Oncol.* 2018;15(5):273–291.
- Cheung LW, Hennessy BT, Li J, Yu S, Myers AP, Djordjevic B, et al. High frequency of PIK3R1 and PIK3R2 mutations in endometrial cancer elucidates a novel mechanism for regulation of PTEN protein stability. *Cancer Discov.* 2011;1(2):170–185.
- Faivre EJ, Lange CA. Progesterone receptors upregulate Wnt-1 to induce epidermal growth factor receptor transactivation and c- Src-dependent sustained activation of Erk1/2 mitogen-activated protein kinase in breast cancer cells. *Mol Cell Biol.* 2007;27(2):466–480.

38. Clemm DL, Sherman L, Boonyaratanakornkit V, Schrader WT, Weigel NL, Edwards DP. Differential hormone-dependent phosphorylation of progesterone receptor A and B forms revealed by a phosphoserine site-specific monoclonal antibody. *Mol Endocrinol*. 2000;14(1):52–65.
39. Liang J, Shang Y. Estrogen and cancer. *Annu Rev Physiol*. 2013;75:225–40.
40. Saxton RA, Sabatini DM. mTOR signaling in growth, metabolism, and disease. *Cell*. 2017;168(6):960–976.
41. Wan X, Harkavy B, Shen N, Grohar P, Helman LJ. Rapamycin induces feedback activation of Akt signaling through an IGF-1R-dependent mechanism. *Oncogene*. 2007;26(13):1932–1940.
42. Salvesen HB, Carter SL, Mannelqvist M, Dutt A, Getz G, Stefansson IM, et al. Integrated genomic profiling of endometrial carcinoma associates aggressive tumors with indicators of PI3 kinase activation. *Proc Natl Acad Sci USA*. 2009;106(12):4834–4839.
43. Slomovitz BM, Coleman RL. The PI3K/AKT/mTOR pathway as a therapeutic target in endometrial cancer. *Clin Cancer Res*. 2012;18(21):5856–5864.
44. Saegusa M, Okayasu I. Frequent impairment of beta-catenin/tcf-4 signaling in both adenoma and carcinoma of the endometrium suggests these are early events in carcinogenesis. *Br J Cancer*. 2001;85(1):81–87.
45. Slater M, Cooper M, Murphy CR. The cytoskeletal proteins alpha-actinin, Ezrin, and talin are De-expressed in endometriosis and endometrioid carcinoma compared with normal uterine epithelium. *Appl Immunohistochem Mol Morphol*. 2007 Jun;15(2):170–4.
46. Byron SA, Gartsman ML, Nelson CM, Dooney CM, Bair RH, Navarro BC, et al. The N550K/H mutations in FGFR2 confer differential resistance to PD173074, dovitinib, and ponatinib ATP-competitive inhibitors. *PLoS One*. 2013;8(9):e54299.
47. Aghajanian C, Sill MW, Darcy KM, Greer B, McMeekin DS, Rose PG, et al. Phase II trial of bevacizumab in recurrent or persistent endometrial cancer: a Gynecologic Oncology Group study. *J Clin Oncol*. 2011;29(16):2259–2265.
48. Santin AD, Bellone S, Van Stedum S, Bushen W, Palmieri M, Siegel ER, et al. Amplification of c-erbB2 oncogene: a major prognostic indicator in uterine serous papillary carcinoma. *Cancer*. 2005;104(7):1391–1397.
49. Ignatov T, Eggemann H, Semczuk A, Smith B, Bischoff J, Roessner A, et al. Role of PTEN expression and ESR1 methylation in endometrial cancer. *J Cancer Res Clin Oncol*. 2010;136(2):223–231.
50. Xiong Y, Dowdy SC, Gonzalez Bosquet J, Zhao Y, Eberhardt NL, Podratz KC, et al. Epigenetic-mediated upregulation of progesterone receptor B gene in endometrial cancer cell lines. *Gynecol Oncol*. 2005;99(1):135–141.
51. Kawabata A, Baojiang C, Bai X, Saeidi N, Barger CJ, Karpf AR, et al. Loss of HDAC-mediated repression and gain of NF-κB activation underlie cytokine induction in ARID1A- and PIK3CA-mutant endometrial cancer cells. *Cancer Res*. 2018;78(14):4017–4030.
52. Kim KH, Roberts CW. Targeting EZH2 in cancer. *Nat Med*. 2016;22(2):128–134.
53. Wu Y, Liu X, Zhou Q, Huang C, Meng X, Xu F, et al. Silent information regulator 1 (SIRT1) ameliorates liver fibrosis via promoting activated stellate cell apoptosis and reversion. *Toxicol Appl Pharmacol*. 2015;289(2):163–176.
54. Huang L, Liao LM, Liu AW, Wu JB, Cheng XL, Lin JX, et al. Overexpression of long noncoding RNA HOTAIR predicts a poor prognosis in patients with cervical cancer. *Arch Gynecol Obstet*. 2014;290(4):717–723.
55. Xie C, Powell C, Yao M, Wu J, Dong Q. Ubiquitin-activating enzyme E1 as a novel therapeutic target for the treatment of restenosis. *Cardiovasc Res*. 2014;101(1):104–113.
56. Burzawa JK, Schmeler KM, Soliman PT, Meyer LA, Bevers MW, Bodurka DC, et al. Prospective evaluation of insulin resistance among endometrial cancer patients. *Am J Obstet Gynecol*. 2011;204(4):355.e1–7.
57. Doll A, Abal M, Rigau M, Monge M, Gonzalez M, Morote J, et al. Novel molecular profiles of endometrial cancer-new light through old windows. *J Steroid Biochem Mol Biol*. 2008;108(3–5):221–229.
58. Huang CY, Chiang SF, Ke TW, Chen TW, You YS, Chen WT, et al. Clinical significance of programmed death-ligand 1 expression in colorectal cancer and its relationship with mucinous differentiation. *Oncotarget*. 2016;7(17):24400–24412.
59. Gottesman MM, Fojo T, Bates SE. Multidrug resistance in cancer: role of ATP-dependent transporters. *Nat Rev Cancer*. 2002;2(1):48–58.
60. Gerlinger M, Rowan AJ, Horswell S, Math M, Larkin J, Endesfelder D, et al. Intratumor heterogeneity and branched evolution revealed by multiregion sequencing. *N Engl J Med*. 2012;366(10):883–892.
61. Stelloo E, Nout RA, Osse EM, Jürgenliemk-Schulz JJ, Jobsen JJ, Lutgens LC, et al. Improved risk assessment by integrating molecular and clinicopathological factors in early-stage endometrial cancer — combined analysis of the PORTEC cohorts. *Clin Cancer Res*. 2016;22(16):4215–4224.
62. Concin N, Matias-Guiu X, Vergote I, Cibula D, Mirza MR, Marnitz S, et al. ESGO/ESTRO/ESP guidelines for the management of patients with endometrial carcinoma. *Int J Gynecol Cancer*. 2021;31(1):12–39.
63. Salvesen HB, Akslen LA. Significance of tumour-associated macrophages, vascular endothelial growth factor and thrombospondin-1 expression for tumour angiogenesis and prognosis in endometrial carcinomas. *Int J Cancer*. 1999;84(5):538–543.
64. Bae GE, Kwon YA, Kim JH, et al. PTEN expression and its association with hormonal therapy response in endometrial carcinoma. *Pathol Res Pract*. 2020;216(11):153188.
65. Bosse T, Peters EE, Creutzberg CL, Jürgenliemk-Schulz IM, Jobsen JJ, Mens JW, et al. Substantial lymph-vascular space invasion (LVSI) is a significant risk factor for recurrence in endometrial cancer, a pooled analysis of PORTEC 1 and 2 trials. *Eur J Cancer*. 2015;51(13):1742–1750.
66. Kim JI, Lim SY, Kim SH, et al. Circulating tumour DNA as a liquid biopsy biomarker in endometrial cancer: a systematic review. *Cancers (Basel)*. 2023;15(4):1245.
67. Oza AM, Elit L, Tsao MS, et al. Phase II study of temsirolimus in women with recurrent or metastatic endometrial cancer. *J Clin Oncol*. 2011;29(24):3278–3285.
68. Slomovitz BM, Jiang Y, Yates MS, Soliman PT, Johnston T, Nowakowski M, et al. Phase II study of everolimus and letrozole in patients with recurrent endometrial carcinoma. *J Clin Oncol*. 2015;33(8):930–936.
69. Soliman PT, Westin SN, Iglesias DA, Fellman BM, Yuan Y, Zhang Q, et al. Everolimus, letrozole, and metformin in women with advanced or recurrent endometrioid endometrial cancer: a multi-center, single arm, phase II study. *Clin Cancer Res*. 2020;26(3):581–587.
70. Brana I, Siu LL. Clinical development of phosphatidylinositol 3-kinase inhibitors for cancer treatment. *BMC Med*. 2012;10:161.
71. Matulonis UA, Shapira-Frommer R, Santin AD, Lisyanskaya AS, Pignata S, Vergote I, et al. Antitumor activity and safety of pembrolizumab in patients with advanced recurrent ovarian cancer: results from the phase II KEYNOTE-100 study. *Ann Oncol*. 2019;30(7):1080–1087.
72. O'Malley DM, Maurer KA, Haag J, et al. Phase II trial of palbociclib and letrozole in endometrial cancer (GOG-3027). *Gynecol Oncol*. 2022;167(Suppl 1):S10.
73. Gennigens C, Jerusalem G, Lapaille L, et al. Recurrent or primary metastatic uterine cancer: current and future treatments. *ESMO Open*. 2022;7(4):100532.
74. Makker V, Colombo N, Casado Herráez A, Santin AD, Colombo A, Miller DS, et al. Lenvatinib plus pembrolizumab for advanced endometrial cancer. *N Engl J Med*. 2022;386(5):437–448.

75. Oaknin A, Tinker AV, Gilbert L, Samouëlian V, Mathews C, Brown J, et al. Clinical activity and safety of the anti-programmed death 1 monoclonal antibody dostarlimab for patients with recurrent or advanced mismatch repair-deficient endometrial cancer: a nonrandomized phase 1 clinical trial. *JAMA Oncol.* 2020;6(11):1766–1772.
76. Lee SM, Moon J, Friedlander M, Dorigo O, Chase DM, Wenzel L, et al. Phase 2 study of anastrozole in women with asymptomatic müllerian cancer. *Cancer.* 2011;117(3):489–497.
77. Fader AN, Roque DM, Siegel E, Buza N, Hui P, Abdelghany O, et al. Randomized phase II trial of carboplatin-paclitaxel versus carboplatin-paclitaxel-trastuzumab in uterine serous carcinomas that overexpress human epidermal growth factor receptor 2/neu. *J Clin Oncol.* 2018;36(20):2044–2051.
78. Meric-Bernstam F, Hamilton EP, Beeram M, et al. Trastuzumab deruxtecan (T-DXd) in patients (pts) with HER2-expressing solid tumors: DESTINY-PanTumor02 (DP-02) interim results. *J Clin Oncol.* 2023;41(Suppl 17):LBA3000.
79. Straining R, Eighmy W. Tazemetostat: EZH2 Inhibitor. *J Adv Pract Oncol.* 2022 Mar;13(2):158-163.
80. Mitsuhashi A, Sato Y, Kiyokawa T, Koshizaka M, Hanaoka H, Shozu M. Phase II study of medroxyprogesterone acetate plus metformin as a fertility-sparing treatment for atypical endometrial hyperplasia and endometrial cancer. *Ann Oncol.* 2016;27(2):262–266.

How to cite this article: Sujavudheen R, Dhanusu P, Chinnapparaj P, Haridoss NK. Resistance Mechanisms to Hormonal Therapy in Endometrial Cancer: A Comprehensive Review. *Indian J Pharm Drug Studies.* 2025; Online First.

Funding: None;

Conflicts of Interest: None Stated