

Ethmoidal encephalocele presenting as spontaneous cerebrospinal fluid rhinorrhea in an adult

Lokesh Kumar¹, Narayanam Anantha Sai Kiran², Kiran Kumar VA², Venkatesh M³,
Luis Rafael Moscote-Salazar⁴, Amit Agrawal⁵

From ¹Assistant Professor, Department of ENT, ²Assistant Professor, Department of Neurosurgery, ³Assistant Professor, Department of Radiology, Narayana Medical College Hospital, Chinthareddypalem, Nellore, Andhra Pradesh, India, ⁴Neurosurgery-Critical Care, Red Latino, Organización Latinoamericana de Trauma y cuidado Neurointensivo, Bogota, Colombia, ⁵Professor, Department of Neurosurgery, Narayana Medical College Hospital, Chinthareddypalem, Nellore, Andhra Pradesh, India

Correspondence to: Dr. Amit Agrawal, Department of Neurosurgery, Narayana Medical College Hospital, Chinthareddypalem, Nellore – 524 003, Andhra Pradesh, India. E-mail: dramitagrawal@gmail.com

Received – 14 August 2018

Initial Review – 30 August 2018

Accepted – 02 October 2018

Spontaneous cerebrospinal fluid (CSF) rhinorrhea is an uncommon presentation of meningoencephalocele and is more common in pediatric population [1-3] than in the adult age group [4-8]. A 48-year-old male patient presented with a complaint of spontaneous watery discharge from the nose for 3 months. It was associated with an occasional headache and a low-grade fever. There was no history of vomiting, seizures, loss of consciousness, blurring of vision, or trauma. Furthermore, there was no history of diabetes or hypertension. His pulse rate was 76/min, blood pressure was 120/78 mmHg, and the temperature was normal. Cardiovascular, chest, per abdomen, and spine examination were normal. Higher mental functions and cranial nerves were normal. There were no motor/sensory deficits or meningeal signs. On prolonged sitting, he was having profuse watery discharge from the left nostril. Routine blood investigations were normal except raised total leukocyte count (14,300 cells/mm³). His magnetic resonance imaging (MRI), brain screening, and computerized tomography (CT) cisternography suggested CSF rhinorrhea on the left side with CSF pooling in the left sphenoidal recess. There was a focal defect in the cribriform plate on the left side (Fig. 1). The patient underwent endoscopic closure of CSF leak with fat, fascia lata, fibrin glue, and middle turbinate pedicle flap (Fig. 2). During surgery, there was a presence of encephalocele in the posterior ethmoidal cells (planum sphenoid region) on the left side. Encephalocele was traced till its origin, cauterized with bipolar and removed. Exploration was done for any other leaks in the posterior ethmoidal region on the right side, but no leak was found. The patient recovered well after surgery and there was no further leak at 6-month follow-up.

Meningoencephalocele is characterized by herniation of brain tissue through a skull defect [9] and basal encephaloceles are an uncommon cause of spontaneous CSF rhinorrhea in adults [10-14]. High-resolution multislice CT scan may not detect the site of the leak as it has low sensitivity; however, an addition of contrast, i.e., CT cisternography increases the sensitivity 80–85% in cases with an active leak [5]. MRI has a higher detection rate in localizing the site of leak including a presence on encephalocele,




Figure 1: Coronal magnetic resonance imaging fast imaging employing steady-state acquisition images (a and b) showing pooling of cerebrospinal fluid (CSF) and herniated left frontal lobe (arrow) into the left ethmoidal air cells suggestive of meningoencephalocele and (c) coronal computerized tomography image shows bony defect (arrow) in the left cribriform plate with pooled CSF density fluid




Figure 2: Endoscopic images (a and b) demonstrating the herniated brain parenchyma covered with meninges (arrow)

and it is complementary to the CT scan (CT provides the bony details for surgical planning) [11,14,15]. Management of ethmoidal encephalocele requires a team approach consisting of an experienced neurosurgeon and otolaryngologists familiar with endoscopic skull base repairs of the skull base defects [3,4,6,11,14,16-18]. Our patient did well after transnasal endoscopic repair of the defect.

REFERENCES

1. Shimizu T, Kitamura S, Kinouchi K, Fukumitsu K. A rare case of upper airway obstruction in an infant caused by basal encephalocele complicating facial midline deformity. *Pediatr Anesth* 1999;9:73-6.
2. Rahbar R, Resto VA, Robson CD, Perez-Atayde AR, Goumnerova LC, McGill TJ, *et al.* Nasal glioma and encephalocele: Diagnosis and

- management. *Laryngoscope* 2003;113:2069-77.
3. Patron V, Roger V, Moreau S, Babin E, Hitier M. State of the art of endoscopic frontal sinus cerebrospinal fluid leak repair. *Eur Ann Otorhinolaryngol Head Neck Dis* 2015;132:347-52.
 4. Pelosi S, Bederson JB, Smouha EE. Cerebrospinal fluid leaks of temporal bone origin: Selection of surgical approach. *Skull Base* 2010;20:253.
 5. Wind JJ, Caputy AJ, Roberti F. Spontaneous encephaloceles of the temporal lobe. *Neurosurg Focus* 2008;25:E11.
 6. Arai A, Mizukawa K, Nishihara M, Fujita A, Hosoda K, Kohmura E. Spontaneous cerebrospinal fluid rhinorrhea associated with a far lateral temporal encephalocele. *Neurol Med Chir* 2010;50:243-5.
 7. Kwon JE, Kim E. Middle fossa approach to a temporo sphenoidal encephalocele. *Neurol Med Chir* 2010;50:434-8.
 8. Papanikolaou V, Bibas A, Ferekidis E, Anagnostopoulou S, Xenellis J. Idiopathic temporal bone encephalocele. *Skull Base* 2007;17:311.
 9. Suwanwela C, Suwanwela N. A morphological classification of sincipital encephalomeningoceles. *J Neurosurg* 1972;36:201-11.
 10. Landreneau FE, Mickey B, Coimbra C. Surgical treatment of cerebrospinal fluid fistulae involving lateral extension of the sphenoid sinus. *Neurosurgery* 1998;42:1101-4.
 11. Goodier M, Lubbe D, Andronikou S, Truter R. Spontaneous lateral sphenoid cerebrospinal fluid fistula: MRI diagnosis. *SA J Radiol* 2012;16:12-13.
 12. Wang J, Bidari S, Inoue K, Yang H, Rhoton Jr A. Extensions of the sphenoid sinus: A new classification. *Neurosurgery* 2010;66:797-816.
 13. Pandey AK. Case report: Anteromedial temporosphenoidal encephalocele with a clinically silent lateral bony defect in the greater wing of the sphenoid. *Indian J Radiol Imaging* 2009;19:311.
 14. Lloyd KM, DelGaudio JM, Hudgins PA. Imaging of skull base cerebrospinal fluid leaks in adults. *Radiology* 2008;248:725-36.
 15. Johnson DS, Brennan P, Toland J, O'dwyer A. Magnetic resonance imaging in the evaluation of cerebrospinal fluid fistulae. *Clin Radiol* 1996;51:837-41.
 16. Hanwate R, Thorawade V, Jagade M, Attakil A, Parelkar K, Pandare M, *et al.* CSF rhinorrhoea with encephalocele through sternberg's canal: Our experience. *Int J Otolaryngol Head Neck Surg* 2014;4:50.
 17. Hammer A, Baer I, Geletneky K, Steiner HH. Cerebrospinal fluid rhinorrhea and seizure caused by temporo-sphenoidal encephalocele. *J Korean Neurosurg Soc* 2015;57:298.
 18. Jones V, Virgin F, Riley K, Woodworth BA. Changing paradigms in frontal sinus cerebrospinal fluid leak repair. *Int Forum Allergy Rhinol* 2012;2:227-32.

Funding: None; Conflict of Interest: None Stated.

How to cite this article: Kumar L, Kiran NAS, Kumar VAK, Venkatesh M, Moscote-Salazar LR, Agrawal A. Ethmoidal encephalocele presenting as spontaneous cerebrospinal fluid rhinorrhea in an adult. *Indian J Case Reports*. 2018;4(5):414-415.

Doi: 10.32677/IJCR.2018.v04.i05.029